



Corrigendum to **“TOSCA – an open-source, finite-volume, large-eddy simulation (LES) environment for wind farm flows” published in Wind Energ. Sci., 9, 297–320, 2024**

Sebastiano Stipa¹, Arjun Ajay¹, Dries Allaerts², and Joshua Brinkerhoff¹

¹School of Engineering, University of British Columbia–Okanagan, Kelowna, Canada

²Aerospace Engineering, TU Delft, Delft, the Netherlands

Correspondence: Sebastiano Stipa (sebstipa@mail.ubc.ca)

Published: 26 February 2024

A mistake was made in the file upload during the file validation process, which led to the four panels at the top of Fig. 3 in Stipa et al. (2024) being repeated in the four bottom panels. That figure is now entirely replaced by Fig. 3 of the present corrigendum.

Acknowledgements. The authors thank Herbert Owen of Barcelona Supercomputing Center (BSC-CNS) for bringing the error to their attention.




Figure 3. Vertical profiles averaged over the last simulation hour of, from top to bottom, velocity magnitude, wind angle, potential temperature and heat flux. The initial lapse rate is (left) 1 K km^{-1} and (right) 10 K km^{-1} . Parameters h_0 and h_1 , obtained by fitting the potential temperature profile with the Rampanelli and Zardi (2004) model, correspond to the capping inversion base and center, respectively.